

Separase (A795) polyclonal antibody

Catalog: AP0374

Host: Rabbit

Reactivity: Human

BackGround:

Separase is a cysteine protease that triggers anaphase in all eukaryotes by participating in separation of sister chromatids during mitosis. Once activated, separase hydrolyzes the SCC1 subunit of cohesin, the chromosomal protein complex responsible for sister chromatid cohesion. Separase and cohesin are highly conserved from yeasts to humans. When the cell is not dividing, separase is prevented from cleaving cohesin through its association with another protein, securin. When anaphase is signaled, the securin is ubiquitinated and hydrolyzed by APC/cyclosome, releasing the active separase. Separase is transiently activated between the two meioses and may also be involved in homolog separation.

Product:

1 mg/ml in Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.2.

Molecular Weight:

~ 230 kDa

Swiss-Prot:

Q14674

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB:1:500~1:1000

Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

Separase (A795) polyclonal antibody detects endogenous levels of Separase protein.

DATA:



Western blot (WB) analysis of Separase (A795) pAb at 1:500 dilution

Lane1:EC9706 whole cell lysate(40ug)

Lane2:HEK293T whole cell lysate(40ug)

Lane3:SGC7901 whole cell lysate(40ug)

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151