

N-CoR (H76) polyclonal antibody

Catalog: BS1462

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

Retinoids are metabolites of vitamin A (retinol) and are believed to represent important signaling molecules during vertebrate development and tissue differentiation. Two families of retinoid receptors have been identified. Retinoic acid receptors (RARs), include RAR α , RAR β and RAR γ , each of which have a high affinity for all-trans retinoic acids and belong to the same class of nuclear transcription factors as thyroid hormone receptors, vitamin D3 receptor and ecdysone receptor. Two cofactors that function to repress transcription, designated SMRT and N-CoR, have been shown to associate with the thyroid receptor and RAR in their unliganded state and are released from them upon ligand binding. The carboxy termini of both proteins contain receptor interacting domains while their amino termini contain two previously undescribed repressor domains. SMRT (silencing mediator for RARs and TRs) is 1495 amino acids in length with an estimated molecular weight of 168 kDa. N-CoR (nuclear receptor corepressor) is a protein 2453 amino acids in length with a relative molecular weight of 270 kDa.

Product:

1 mg/ml in Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2.

Molecular Weight:

~ 270 kDa

Swiss-Prot:

O75376

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200

IF: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

NCoR1 (H76) polyclonal antibody detects endogenous levels of NCoR1 protein.

DATA:




Western blot (WB) analysis of N-CoR (H76) pAb at 1:500 dilution

Lane1:CT26 whole cell lysate(40ug)

Lane2:C6 whole cell lysate(40ug)

Lane3:HepG2 whole cell lysate(40ug)

Lane4:Hela whole cell lysate(40ug)



Immunohistochemistry (IHC) analyzes of N-CoR (H76) pAb in paraffin-embedded human breast carcinoma tissue at 1:100.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151



PRODUCT DATA SHEET

Bioworld Technology, Inc.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park,
MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046,
P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151