

**NFAT4 (E159) polyclonal antibody**

Catalog: BS1559

Host: Rabbit

Reactivity: Human, Mouse, Rat

**BackGround:**

NFATc4 (nuclear factor of activated T cells, cytoplasmic, calcineurin-dependent 4) is a member of the nuclear factors of activated T cells DNA-binding transcription complex that influences cytokine gene expression, cardiac hypertrophy and adipocyte differentiation. This complex consists of at least two components, a cytosolic component that translocates to the nucleus upon T cell receptor (TCR) stimulation and an inducible nuclear component. Other members of this family participate in the formation of this complex. NFATc4 plays a role in the inducible expression of cytokine genes in T cells, including the induction of IL-2 and IL-4. p38 MAP kinase phosphorylates multiple residues in the NFAT homology domain of NFATc4.

**Product:**

1 mg/ml in Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2.

**Molecular Weight:**

~ 115 kDa

**Swiss-Prot:**

Q12968

**Purification&Purity:**

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

**Applications:**

IHC: 1:50~1:200

**Storage&Stability:**

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

**Specificity:**

NFAT4 (E159) polyclonal antibody detects endogenous levels of NFAT4 protein.

**DATA:****Note:**

For research use only, not for use in diagnostic procedure.

**Bioworld Technology, Inc.**

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: [info@bioworld.com](mailto:info@bioworld.com)

Tel: 6123263284

Fax: 6122933841

**Bioworld technology, co. Ltd.**

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: [info@biogol.com](mailto:info@biogol.com)

Tel: 0086-025-68037686

Fax: 0086-025-68035151