

## ZAP70 (T286) polyclonal antibody

Catalog: BS1610

Host: Rabbit

Reactivity: Human,Rat

### BackGround:

The activation of T lymphocytes by antigens is mediated by the T cell receptor (TCR), which is a multisubunit complex assembled from at least six different genes . The TCR subunits include the  $\alpha$  and  $\beta$  chains, the CD3  $\gamma$ ,  $\delta$  and  $\epsilon$  chains and a  $\zeta$ -containing homodimer or heterodimer . The protein tyrosine kinase ZAP-70 binds to the phosphorylated immunoreceptor tyrosine-base activation motifs (ITAMs) of the TCR  $\zeta$  chain through two src-homology (SH2) domains . This binding results in the phosphorylation of ZAP-70 on multiple tyrosine residues, including Tyr292 and Tyr319 . ZAP-70 is autophosphorylated on Tyr292, which is thought to negatively regulate ZAP-70 function in lymphocytes . Alternatively, ZAP-70 is positively regulated by phosphorylation on Tyr319, which mediates the SH2-dependent interaction between Lck and ZAP-70 .

### Product:

1 mg/ml in Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2.

### Molecular Weight:

~ 70 kDa

### Swiss-Prot:

P43403

### Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

### Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200


### Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

### Specificity:

ZAP-70 (T286) polyclonal antibody detects endogenous levels of ZAP-70 protein.

### DATA:



Western blot (WB) analysis of ZAP70 (T286) at 1:500 dilution

Lane1:THP-1 whole cell lysate(40ug)

Lane2:Myla2059 whole cell lysate(40ug)

Lane3:HuT78 whole cell lysate(40ug)

Lane4:The Thymus tissue lysate of Rat(40ug)

### Note:

For research use only, not for use in diagnostic procedure.

### Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA.

Email: [info@bioworld.com](mailto:info@bioworld.com)

Tel: 6123263284

Fax: 6122933841

### Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: [info@biogot.com](mailto:info@biogot.com)

Tel: 0086-025-68037686

Fax: 0086-025-68035151