

mTOR (S2442) polyclonal antibody

Catalog: BS1844

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

mTOR, or FKBP12-rapamycin associated protein (FRAP), is one of a family of proteins involved in cell cycle progression, DNA recombination, and DNA damage detection. In rat, it is a 245-kDa protein (symbolized RAFT1) with significant homology to the *Saccharomyces cerevisiae* protein TOR1 and has been shown to associate with the immunophilin FKBP12 in a rapamycin-dependent fashion. The FKBP12-rapamycin complex is known to inhibit progression through the G1 cell cycle stage by interfering with mitogenic signaling pathways involved in G1 progression in several cell types, as well as in yeast. The binding of FRAP to FKBP12-rapamycin correlated with the ability of these ligands to inhibit cell cycle progression.

Product:

1 mg/ml in Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2.

Molecular Weight:

~ 289 kDa

Swiss-Prot:

P42345

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200

IP: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

mTOR (S2442) polyclonal antibody detects endogenous levels of mTOR protein.

DATA:



Western blot (WB) analysis of mTOR (S2442) at 1:500 dilution

Lane1:NIH-3T3 whole cell lysate(40µg)


Lane2:HEK293T whole cell lysate(40µg)

Lane3:H9C2 whole cell lysate(40µg)


Lane4:Hela whole cell lysate(40µg)

Lane5:MCF-7 whole cell lysate(40µg)

Lane6:HepG2 whole cell lysate(40µg)



Immunohistochemistry (IHC) analyzes of mTOR (S2442) pAb in paraffin-embedded human colorectal carcinoma tissue at 1:50.



Immunoprecipitation of SGC7901 cell lysates using mTOR (S2442)

pAb (Sepharose Bead Conjugate)#BD0048 (lane 2 and lane 3) and

Nonspecific IgG Control (Sepharose Bead Conjugate)#BD0048 (lane

4 and lane 5). Lane 1 is 30% input. The western blot was probed using

mTOR (S2442) pAb #BS1844. “↑”(supernatant); “↓”(deposition).

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151



PRODUCT DATA SHEET

Bioworld Technology, Inc.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park,
MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046,
P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151