

Catenin- β (F660) polyclonal antibody

Catalog: BS1982

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

β -Catenin is a key downstream effector in the Wnt signaling pathway. It is implicated in two major biological processes in vertebrates: early embryonic development and tumorigenesis. CK1 phosphorylates β -catenin at Ser45. This phosphorylation event primes β -catenin for subsequent phosphorylation by GSK-3 β . GSK-3 β destabilizes β -catenin by phosphorylating it at Ser33, Ser37, and Thr41. Mutations at these sites result in the stabilization of β -catenin protein levels and have been found in many tumor cell lines.

Product:

1 mg/ml in Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.2.

Molecular Weight:

~ 92 kDa

Swiss-Prot:

P35222

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

IHC: 1:50~1:200

IF: 1:50~1:200

Storage&Stability:



Store at 4 °C short term. Aliquot and store at -20 °C long

term. Avoid freeze-thaw cycles.

Specificity:

Catenin- β (F660) polyclonal antibody detects endogenous levels of Catenin- β protein.

DATA:



Immunohistochemistry (IHC) analyzes of Catenin- β (F660) pAb in paraffin-embedded human breast carcinoma tissue at 1:100.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151