

TANK (F194) polyclonal antibody

Catalog: BS2231

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

The tumor necrosis factor (TNF) receptor superfamily is composed of several type I integral membrane glycoproteins that exhibit homology in their cysteine-rich extracellular domains. Members of this family include TNF-RI and -RII, FAS, OX40, CD27, CD30 and CD40. Ligands for these receptors can be small, secreted proteins such as TNF, or type II integral membrane proteins, such as the CD40 ligand, CD40L. While the signal transduction mechanism of the TNF receptor superfamily is poorly understood, stimulation of cells with either TNF or soluble CD40L has been shown to induce the nuclear translocation of NFκB. Members of the TRAF family associate with activated TNF-R and CD40 and have been implicated in this process. The discovery of a novel protein, designated TANK, has shed light on the means by which TRAF activation of NFκB occurs. TANK is not only capable of binding to all three TRAFs, but also of synergizing with TRAF2 to activate the NFκB signaling cascade. TANK contains a regulatory carboxy terminal domain that maintains its inactivity in unstimulated cells. Upon its association with TRAF2, the inhibitory effect of this domain is overcome.

Product:

1 mg/ml in Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.2.

Molecular Weight:

~48 kDa

Swiss-Prot:

Q92844

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200

IF: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.


Specificity:

TANK (F194) polyclonal antibody detects endogenous levels of TANK protein.

DATA:



Western blot (WB) analysis of TANK (F194) polyclonal antibody in extracts from raw264.7 cells.



Immunohistochemistry (IHC) analyzes of TANK (F194) pAb in paraffin-embedded human kidney carcinoma tissue at 1:50, showing cytoplasmic staining. Negative control (the right) Using PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin followed by avidin-peroxidase.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151



PRODUCT DATA SHEET

Bioworld Technology, Inc.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park,
MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046,
P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151