

Csk (N63) polyclonal antibody

Catalog: BS2275 Host: Rabbit	Reactivity: Human, Mouse, Rat
BackGround: C terminal Src kinase (CSK) is a non receptor protein ty- rosine kinase. One of its major functions is to specifically phosphorylate a conserved C terminal tyrosine of Src family kinases thereby down regulating their kinase ac- tivity. CSK is expressed ubiquitously as a protein of ap- proximately 50 KDa in cytosol, but a fraction distributes to lipid rafts due to interaction with the lipid raft associ- ated protein PAG (CSK binding protein).	 IHC: 1:50~1:200 IF: 1:50~1:200 Storage&Stability: Store at 4 ℃ short term. Aliquot and store at -20 ℃ long term. Avoid freeze-thaw cycles. Specificity: Csk (N63) polyclonal antibody detects endogenous levels of Csk protein. DATA:
Product: 1 mg/ml in Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.2. Molecular Weight: ~ 50 kDa Swiss-Prot: P41240	120kd 90kd 50kd Csk (N63) 34kd HepG2 whole cell lysate Csk (N63) pAb at 1:500 dilution <i>By linwald Tech</i>
Purification&Purity: The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific im- munogen and the purity is > 95% (by SDS-PAGE). Applications:	Note: For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc. Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA. Email: info@bioworlde.com Tel: 6123263284 Fax: 6122933841

Bioworld technology, co. Ltd. Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China. Email: <u>info@biogot.com</u> Tel: 0086-025-68037686 Fax: 0086-025-68035151