



Nogo A (E499) polyclonal antibody

Catalog: BS2888

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

CNS white matter is selectively inhibitory for axonal out-growth. Nogo (also designated NI250 and Reticulon 4-A) is an oligodendrocyte-specific member of the Reticulon family and is a component of CNS white matter that inhibits axon outgrowth, induces collapse of growth cones of chick dorsal root ganglion cells, and inhibits the spreading of 3T3 fibroblasts. Other members of the Reticulon protein family do not inhibit axon extension and are thought to have a role in ER function. Nogo is expressed by oligodendrocytes but not by Schwann cells, and associates primarily with the endoplasmic reticulum. Nogo exists in three different splice forms, Nogo A, Nogo B and Nogo C. Nogo A and Nogo C have a molecular mass of 200-250 kDa and 22 kDa, respectively.

Product:

1 mg/ml in Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2.

Molecular Weight:

~ 130 kDa

Swiss-Prot:

Q9NQC3

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200

Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

Nogo A (E499) polyclonal antibody detects endogenous levels of Nogo A protein.

DATA:

Immunohistochemistry (IHC) analyzes of Nogo A (E499) pAb in paraffin-embedded human skeletal muscle tissue.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogol.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151