

CMTM3 (E13) polyclonal antibody

Catalog:	BS3285	Host:	Rabbit	Reactivity: Human, Mouse, Rat		
BackGround:				Specificity:		
CMTM3 belongs to the chemokine-like factor gene su-			tor gene su-	CMTM3(E13) pAb detects endogenous levels of		
perfamily, a novel family that is similar to the chemokine			e chemokine	CMTM3 protein.		
and the transmembrane 4 superfamilies of signaling mol-				DATA:		
ecules. This gene is one of several chemokine-like factor genes located in a cluster on chromosome 16.				1 2 3 4 5 130 95 72 55		
Product:				43		
1 mg/ml in Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.3.			with 0.05%	34 26 CMTM 3		
Molecular Weight:						
~ 20 kDa				ZX180821-C3		
Swiss-Prot:				Western blot (WB) analysis of CMTM3 (E13) pAb at 1:500 dilution		
Q96MX0				Lane1:K562 whole cell lysate(40ug)		
Purification&Purity:				Lane2:A375 whole cell lysate(40ug)		
The antibody was affinity-purified from rabbit antiserum			oit antiserum	Lane3:Hela whole cell lysate(40ug)		
by affinity	by affinity-chromatography using epitope-specific im-			Lane4:The Embryo tissue lysate of Mouse(40ug)		
munogen and the purity is > 95% (by SDS-PAGE). Applications: WB: 1:1000~1:2000 Storage&Stability:			AGE).	Lane5:The Testis tissue lysate of Rat(40ug)		
				Note:		
				For research use only, not for use in diagnostic procedure.		
Store at 4 %	C short term. Alic	uot and store a	t -20 ℃ long	-		
term. Avoid	freeze-thaw cycle	es.				

Bioworld Technology, Inc. Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA. Email: <u>info@bioworlde.com</u> Tel: 6123263284 Fax: 6122933841

Bioworld technology, co. Ltd. Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China. Email: <u>info@biogot.com</u> Tel: 0086-025-68037686 Fax: 0086-025-68035151