

## MNDA (V406) polyclonal antibody

Catalog: BS3332

Host: Rabbit

Reactivity: Human, Mouse, Rat

### BackGround:

Interferon-inducible proteins include IFI-202, IFI-203, IFI-204 and D3, which are encoded by six or more structurally related and IFN-inducible mouse genes mapping at the q21-q23 region of chromosome 1. The proteins encoded by these genes have homologous 200 amino acid segments. IFI-202 is a 52 kDa, primarily nuclear phosphoprotein which inhibits cell growth, in part by modulating transcriptional activity of NFκB, E2F, AP-1 and p53. Two related human proteins, MNDA (myeloid cell nuclear differentiation antigen) and IFI-16, have also been described. Expression of MNDA has been observed specifically in cells of the granulocyte-macrophage lineage. IFI-16 is constitutively expressed in various T and B cell lines and can be induced by IFN-γ in HL60 cells. At least four of the Gene 200 cluster of IFN-inducible proteins, IFI-202, IFI-204, MNDA and IFI-16, are localized in the nucleus.

### Product:

1 mg/ml in Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2.

### Molecular Weight:

~ 46 kDa

### Swiss-Prot:

P41218

### Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

### Applications:

IHC: 1:50~1:200


### Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.


### Specificity:

MNDA (V406) polyclonal antibody detects endogenous levels of MNDA protein.

### DATA:



Immunohistochemistry (IHC) analyzes of MNDA (V406) pAb in paraffin-embedded human tonsil carcinoma tissue at 1:50. showing cytoplasmic and nucleus staining. Negative control (the right) Using PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin followed by avidin-peroxidase.



Immunohistochemistry (IHC) analyzes of MNDA (V406) pAb in paraffin-embedded human kidney carcinoma tissue at 1:50. showing cytoplasmic and nucleus staining. Negative control (the right) Using PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin followed by avidin-peroxidase.

### Note:

For research use only, not for use in diagnostic procedure.

### Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: [info@bioworld.com](mailto:info@bioworld.com)

Tel: 6123263284

Fax: 6122933841

### Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: [info@biogot.com](mailto:info@biogot.com)

Tel: 0086-025-68037686

Fax: 0086-025-68035151