

PRODUCT DATA SHEET

Bioworld Technology, Inc.

ARHGAP17 (T363) polyclonal antibody

Catalog: BS3390 Host: Rabbit Reactivity: Human, Mouse, Rat

BackGround:

GTPase-activating proteins (GAPs) accelerate the intrinsic rate of GTP hydrolysis of Ras-related proteins, resulting in down regulation of their active form. ARHGAP17 (Rho GTPase activating protein 17), also known as RICH1, WBP15, MST066, MST110, NADRIN, PP4534, RICH1B, MSTP038, MSTP066 or MSTP110, is a ubiquitously expressed peripheral membrane protein whose expression is highest in heart and placenta. ARHGAP17 is involved in the maintenance of tight junctions by regulating the activity of Cdc42, thereby playing a central role in apical polarity of epithelial cells. Containina a BAR domain and a Rho-GAP domain, ARHGAP17 acts as a GTPase activator for the Cdc42 GTPase by converting it to an inactive GDP-bound state. ARHGAP17 may also participate in the Ca2+-dependent regulation of exocytosis by catalyzing

GTPase activity of Rho family proteins and by inducing the reorganization of cortical actin filaments. ARHGAP17 exists as seven alternative splice variants.

Product:

1 mg/ml in Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.3.

Molecular Weight:

~ 130 kDa

Swiss-Prot:

Q68EM7

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000


Storage&Stability:

Store at $4 \, \mathbb{C}$ short term. Aliquot and store at $-20 \, \mathbb{C}$ long term. Avoid freeze-thaw cycles.

Specificity:

ARHGAP17(T363) pAb detects endogenous levels of ARHGAP17 protein.

DATA:



Western blot (WB) analysis of ARHGAP17 (T363) pAb at 1:1000 dilu-

Lane1:Hela whole cell lysate(30ug)

Lane2:HCT116 whole cell lysate(40ug)

Lane3:A375 whole cell lysate(40ug)

Lane4:The Embryo tissue lysate of Mouse(40ug)

Lane5:H9C2 whole cell lysate(40ug)

Note

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park,

MN 55416,USA.

Email: <u>info@bioworlde.com</u>

Tel: 6123263284 Fax: 6122933841 Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046,

P. R. China.

Email: <u>info@biogot.com</u> Tel: 0086-025-68037686 Fax: 0086-025-68035151