

## WDHD1 (Q756) polyclonal antibody

Catalog:	BS3414	Host:	Rabbit	Reactivity: Human
BackGround:				by affinity-chromatography using epitope-specific im-
The protein encoded by this gene contains multiple				munogen and the purity is $> 95\%$ (by SDS-PAGE).
N-terminal WD40 domains and a C-terminal high mobil-				Applications:
ity group (HMG) box. WD40 domains are found in a va-				WB: 1:500~1:1000
riety of eukaryotic proteins and may function as adap-				IHC: 1:50~1:200
tor/regulatory modules in signal transduction, pre-mRNA				Storage&Stability:
processing and cytoskeleton assembly. HMG boxes are				Store at 4 °C short term. Aliquot and store at -20 °C long
found in many eukaryotic proteins involved in chromatin				term. Avoid freeze-thaw cycles.
assembly, transcription and replication. Alternative splic-			rnative splic-	Specificity:
ing results in two transcript variants encoding different			ing different	WDHD1 (Q756) polyclonal antibody detects endogenous
isoforms.				levels of WDHD1 protein.
Product:				DATA:
1 mg/ml in Phosphate buffered saline (PBS) with 0.05%				
sodium azide, approx. pH 7.2. Molecular Weight:				Immunohistochemistry (IHC) analyzes of WDHD1 (Q756) pAb in par-
				affin-embedded human breast cancer tissue.
~ 126 kDa				Note:
Swiss-Prot	t:			For research use only, not for use in diagnostic procedure.
075717				

**Purification&Purity:** 

The antibody was affinity-purified from rabbit antiserum