

SDF-1 (M93) polyclonal antibody

Catalog: BS3512

Host: Rabbit

Reactivity: Human, Mouse, Rat

Background:

SDF-1 was originally described as a pre-B cell stimulatory factor, but has now been shown to function as a potent chemoattractant for T cells and monocytes, but not neutrophils. Receptors for the C-X-C family are G protein-coupled, seven-pass, transmembrane domain proteins which include IL-8RA, IL-8RB and fusin (also designated LESTR or CXCR-4). Fusin is highly homologous to the IL-8 receptors, sharing 37% sequence identity at the amino acid level. The IL-8 receptors bind to IL-8, NAP-2 and MSGA, while fusin binds to its cognate ligand, SDF-1. Fusin has been identified as the major co-receptor for T-tropic HIV-1, and SDF-1 has been shown to inhibit HIV-1 infection. Six human SDF-1 isoforms exist due to alternative splicing of CXCL12, the gene encoding SDF-1.

Product:

1 mg/ml in Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2.

Molecular Weight:

~ 10 kDa

Swiss-Prot:

P48061

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific im-

munogen and the purity is > 95% (by SDS-PAGE).

Applications:

IHC:1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

SDF-1 (M93) polyclonal antibody detects endogenous levels of SDF-1 protein.

DATA:



Immunohistochemistry (IHC) analyzes of SDF-1 (M93) pAb in paraffin-embedded human brain tissue.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151