



Trypsin-3 (M166) polyclonal antibody

Catalog: BS3707

Host: Rabbit

Reactivity: Human

BackGround:

In the small intestine, each isoform is cleaved by Enterokinase into its active form, Trypsin-1, Trypsin-2 and Trypsin-3, respectively. All trypsins are members of the serine protease trypsin family. The activated trypsins go on to activate other protease zymogens and play a role in the autoactivation of trypsinogens. This suggests an important role for trypsins in digestion. Mutations in the gene encoding Trypsin-1 that stimulate its activity are associated with autosomal dominant hereditary pancreatitis (HCP), also known as chronic pancreatitis (CP), a disease characterized by persistent, severe abdominal pain due to calcifications of the parenchyma, pancreatic stones, cysts and pancreatic head enlargement. Trypsin-3 is expressed in the brain in addition to the pancreas.

Product:

1 mg/ml in Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2.

Molecular Weight:

~ 32 kDa

Swiss-Prot:

P35030

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

IHC: 1:50~1:200

Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

Trypsin-3 (M166) polyclonal antibody detects endogenous levels of Trypsin-3 protein.

DATA:

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogol.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151