

SLP-76 (G122) polyclonal antibody

Catalog: BS3758

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

Overexpression of Vav or SLP-76 in Jurkat cells leads to NFAT activation and IL-2 production. When co-expressed, Vav and SLP-76 synergize to induce a robust basal and TCR-mediated IL-2 response. Although SLP-76 does not contain a motif that would indicate it to be a member of the tyrosine, serine/threonine or lipid kinase families, it does contain several putative SH2/SH3-binding domains and has been shown to physically associate with the adapter protein GRB2 as well as PLC g1. The discovery of SLP-76 represents an important step in elucidating the mechanism of Vav transformation and TCR-mediated NFAT activation.

Product:

1 mg/ml in Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2.

Molecular Weight:

~ 75 kDa

Swiss-Prot:

Q13094

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

SLP-76 (G122) polyclonal antibody detects endogenous levels of SLP-76 protein.

DATA:



Immunohistochemistry (IHC) analyzes of SLP-76 (G122) pAb in paraffin-embedded human breast carcinoma tissue at 1:100.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151