

Histone H3 (mono+di+tri methyl K79) Rabbit monoclonal anti-body

Catalog: BS40149

Host: Rabbit

Reactivity: Human Mouse Rat

BackGround:

H3 Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling. The nucleosome is a histone octamer containing two molecules each of H2A, H2B, H3 and H4 assembled in one H3-H4 heterotetramer and two H2A-H2B heterodimers.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2.

Molecular Weight:

15kDa

Swiss-Prot:

P68431

Purification&Purity:

Affinity-chromatography

Applications:

WB 1:500~1:1000 IHC 1:1000~1:2000


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.


Specificity:

Histone H3 (mono+di+tri methyl K79) Antibody detects endogenous levels of total Histone H3 (mono+di+tri methyl K79)

DATA:



Western blot analysis of Histone H3 (mono+di+tri methyl K79) expression in (1) NIH/3T3 cell lysate; (2) A549 cell lysate.



Immunohistochemical analysis of paraffin-embedded rat colon, using Histone H3 (mono+di+tri methyl K79) Antibody.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151