

## Inhibin alpha Rabbit monoclonal antibody

Catalog:	BS41039	Host:	Rabbit	Reactivity:	Human
BackGround:				WB 1:500~1:1000 IHC 1:50~1:200	
Inhibins and activins inhibit and activate, respectively, the				Storage&Stability:	
secretion of follitropin by the pituitary gland. In-			gland. In-	Store at 4 ${\rm C}$ short term. Aliquot and store at -20 ${\rm C}$ long	
hibins/activins are involved in regulating a number of di-				term. Avoid freeze-thaw cycles.	
verse functions such as hypothalamic and pituitary hor- mone secretion, gonadal hormone secretion, germ cell development and maturation, erythroid differentiation,			oituitary hor-	Specificity:	
			n, germ cell	Inhibin alpha Antibody detects endogenous levels of total Inhibin alpha	
			fferentiation,		
insulin sec	nsulin secretion, nerve cell survival, embryonic axial development or bone growth, depending on their subunit			DATA:	
-				KDa   250   1300   100   75   50   37   25   20   150   150	
composition. <b>Product:</b> Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2.					
			odium azide,		
Molecular Weight:					
40kDa	40kDa				
Swiss-Prot	Swiss-Prot:			Western blot analysis of Inhibin alpha expression in Human testis lysate.	
P05111	P05111			Note:	
Purification&Purity:				For research use only, not for use in diagnostic procedure.	
Affinity-chromatography					
Applications:					