

p53 (phospho-S9) polyclonal antibody

Catalog: BS4149

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

p53 plays a major role in the cellular response to DNA damage and other genomic aberrations. The activation of p53 can lead to either cell cycle arrest and DNA repair, or apoptosis. p53 is phosphorylated at multiple sites in vivo and by several different protein kinases in vitro. p53 can apparently be phosphorylated by ATM, ATR, and DNAPK at Ser15; the phosphorylation impairs the ability of MDM2 to bind p53, promoting both the accumulation and functional activation of p53 in response to DNA damage. p53 is phosphorylated at Ser6 and Ser9 by ck1d and ck1e both in vitro and in vivo. Phosphorylation of p53 at Ser46 is important in regulating the ability of p53 at Ser33 by cdk7/cyclin H and in response to UV irradiation has been observed.

Product:

1 mg/ml in Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2.

Molecular Weight:

~ 53,43 kDa

Swiss-Prot:

P04637

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

IHC: 1:50~1:200

Storage&Stability:

Store at $4 \,^{\circ}{\rm C}$ short term. Aliquot and store at $-20 \,^{\circ}{\rm C}$ long term. Avoid freeze-thaw cycles.

Specificity:

p-p53 (S9) polyclonal antibody detects endogenous levels of p53 protein when phosphorylated at Ser9.

DATA:



Immunohistochemistry (IHC) analyzes of p-p53 (S9) pAb in paraf-

fin-embedded human breast carcinoma tissue at 1:100.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

 Add:
 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA.

 Email:
 info@bioworlde.com

 Tel:
 6123263284

 Fax:
 6122933841

Bioworld technology, co. Ltd. Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China. Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151