

Phospho-beta Catenin (Thr41/Ser45) Recombinant Rabbit mAb

adenomatous polyposis of the colon. Mutations in this gene are a cause of colorectal cancer (CRC), pilomatrix- oma (PTR), medulloblastoma (MDB), and ovarian cancer. Alternative splicing results in multiple transcript variants.	Catalog:	BS43345	Host:	Rabbit	Reactivity:	Human
binds to the product of the APC gene, which is mutated in adenomatous polyposis of the colon. Mutations in this gene are a cause of colorectal cancer (CRC), pilomatrix- oma (PTR), medulloblastoma (MDB), and ovarian cancer. Alternative splicing results in multiple transcript variants.	BackGroun The protein proteins that necessary for cell layers tween cells. cytoskeletor contact inhi	BackGround: The protein encoded by this gene is part of a complex of proteins that constitute adherens junctions (AJs). AJs are necessary for the creation and maintenance of epithelial cell layers by regulating cell growth and adhesion be- tween cells. The encoded protein also anchors the actin cytoskeleton and may be responsible for transmitting the contact inhibition signal that causes cells to stop dividing			Molecular Weight: 86 kDa Swiss-Prot: P35222 Purification&Purity: Affinity Purification Applications:	2
[provided by Kelseq, Aug 2010]	binds to the product of the APC gene, which is mutated in adenomatous polyposis of the colon. Mutations in this gene are a cause of colorectal cancer (CRC), pilomatrix- oma (PTR), medulloblastoma (MDB), and ovarian cancer.				Storage&Stability: Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles. Isotype: IgG	

Product:

Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.

Note:

For research use only, not for use in diagnostic procedure.