

Crotonyl-Histone H4 (Lys8) Recombinant Rabbit mAb

BackGround:P62805Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a tem- plate. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromoso- mal stability. DNA accessibility is regulated via a com- plex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.Pdification&Purity:Product:Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.Note: For research use only, not for use in diagnostic procedure.Note:Swiss-Prot:	Catalog:	BS43432	Host:	Rabbit	Reactivity: Human, Mouse, Rat
plex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.term. Avoid freeze-thaw cycles.Product:Isotype:Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.DATA:Molecular Weight:Note:11 kDaFor research use only, not for use in diagnostic procedure.	BackGround:Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a tem- plate. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromoso- mal stability. DNA accessibility is regulated via a com- plex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.Product:Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.Molecular Weight:			es wrap and accessibility A as a tem- ranscription I chromoso-	P62805 Purification&Purity: Affinity Purification Applications: WB: 1:1000 IHC: 1:100 ICC/IF: 1:200 Storage&Stability:
Product: Isotype: Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), IgG 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and DATA: 0.05% BSA. Stable for 12 months from date of receipt. Note: Molecular Weight: For research use only, not for use in diagnostic procedure. 11 kDa Image: Note:					
Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), DATA: 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and DATA: 0.05% BSA. Stable for 12 months from date of receipt. Note: Molecular Weight: For research use only, not for use in diagnostic procedure. 11 kDa Note:				odeling.	
0.05% BSA. Stable for 12 months from date of receipt. Note: Molecular Weight: For research use only, not for use in diagnostic procedure. 11 kDa				-	
11 kDa					
Swiss-Prot:					For research use only, not for use in diagnostic procedure.
	Swiss-Prot	:			