

PKA α / β cat (phospho-T197) polyclonal antibody

Catalog: BS4345

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

PRKACA and PRKACB are members of the Ser/Thr protein kinase family and are a catalytic subunit of cAMP-dependent protein kinase. cAMP is a signaling molecule important for a variety of cellular functions. cAMP exerts its effects by activating the cAMP-dependent protein kinase, which transduces the signal through phosphorylation of different target proteins. The inactive kinase holoenzyme is a tetramer composed of two regulatory and two catalytic subunits. cAMP causes the dissociation of the inactive holoenzyme into a dimer of regulatory subunits bound to four cAMP and two free monomeric catalytic subunits.

Product:

1 mg/ml in Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.2.

Molecular Weight:

~ 42 kDa

Swiss-Prot:

P17612/P22694

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200

Storage&Stability:


Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

p-PKA α / β cat (T197) polyclonal antibody detects endog-

enous levels of PKA α / β cat protein when phosphorylated at Thr197.

DATA:




Western blot (WB) analysis of P-PKA α / β cat (T197) pAb at 1:500 dilution

Lane1: HEK293T whole cell lysate(40ug)

Lane2: A549 whole cell lysate(40ug)

Lane3: The Testis tissue lysate of Mouse(40ug)

Lane4: The Testis tissue lysate of Rat(40ug)



Immunohistochemistry (IHC) analyzes of p-PKA α / β cat (T197) pAb in paraffin-embedded human colorectal carcinoma tissue at 1:50.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151