

SNAI1 Recombinant Rabbit mAb

BackGround: WB: 1:1000 >IP:1:20 The Drosophila embryonic protein snail is a zinc finger transcriptional repressor which downregulates the expression of ectodermal genes within the mesodern. The nuclear protein encoded by this gene is structurally similar to the Drosophila snail protein, and is also thought to be critical for mesoderm formation in the developing embryo. At least two variants of a similar processed pseudogene have been found on chromosome 2. [provided by RefSeq, Jul 2008] Isotype: Product: DATA: Store at -20°C. Supplied in 50mM Tris-Glycine(PH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt. Western blot analysis of extracts from HepG2.SW620 cells using BS48474 at 1: 1000. Swiss-Prot: O95863 For research use only, not for use in diagnostic procedure on the meson of the meson o	Catalog:	BS48474	Host:	Rabbit	Reactivity:	Human
Storage&Stability:Storage&Stability:storage<	BackGrou	nd			WB: 1:1000 IP: 1	:20
transcriptional repressor which downregulates the expression of ectodermal genes within the mesoderm. The nuclear protein encoded by this gene is structurally similar to the Drosophila snail protein, and is also thought to be critical for mesoderm formation in the developing em- bryo. At least two variants of a similar processed pseudo- gene have been found on chromosome 2. [provided by RefSeq, Jul 2008]Isotype: IgGProduct: Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.Western blot analysis of extracts from HepG2,SW620 cells using BS48474 at 1: 1000.Swiss-Prot: O95863 Purification&Purity: Affinity PurificationNote: For research use only, not for use in diagnostic procedure					Storage&Stability:	
sion of ectodermal genes within the mesoderm. The nuclear protein encoded by this gene is structurally similar to the Drosophila snail protein, and is also thought to be critical for mesoderm formation in the developing embryo. At least two variants of a similar processed pseudogene have been found on chromosome 2. [provided by RefSeq, Jul 2008] Product: Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt. Molecular Weight: 29 kDa Stwiss-Prot: O95863 Purification&Purity: Affinity Purification					0	n. Aliquot and store at -20 °C long
clear protein encoded by this gene is structurally similar to the Drosophila snail protein, and is also thought to be critical for mesoderm formation in the developing em- bryo. At least two variants of a similar processed pseudo- gene have been found on chromosome 2. [provided by RefSeq, Jul 2008]Isotype: IgGProduct:DATA:Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.Western blot analysis of extracts from HepG2,SW620 cells using BS48474 at 1: 1000.Stwiss-Prot:Note:O95863For research use only, not for use in diagnostic procedure For research use only, not for use in diagnostic procedure						
to the Drosophila snail protein, and is also thought to be critical for mesoderm formation in the developing em- bryo. At least two variants of a similar processed pseudo- gene have been found on chromosome 2. [provided by RefSeq, Jul 2008]IgGProduct:DATA:Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.Western blot analysis of extracts from HepG2,SW620 cells using BS48474 at 1: 1000.Swiss-Prot:Note:O95863For research use only, not for use in diagnostic procedure For research use only, not for use in diagnostic procedureAffinity PurificationHermite and Comparison						5
critical for mesoderm formation in the developing embryo. At least two variants of a similar processed pseudo- gene have been found on chromosome 2. [provided by RefSeq, Jul 2008]DATA:Product:DATA:Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.Western blot analysis of extracts from HepG2,SW620 cells using BS48474 at 1: 1000.Swiss-Prot:Note:O95863For research use only, not for use in diagnostic procedure For research use only, not for use in diagnostic procedure					**	
gene have been found on chromosome 2. [provided by RefSeq, Jul 2008] Product: Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt. Molecular Weight: 29 kDa Swiss-Prot: O95863 Purification&Purity: Affinity Purification	critical for mesoderm formation in the developing em-					
RefSeq, Jul 2008]Product:Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.Molecular Weight:Western blot analysis of extracts from HepG2,SW620 cells using BS48474 at 1: 1000.Swiss-Prot:Note:O95863For research use only, not for use in diagnostic procedure For research use only, not for use in diagnostic procedure Affinity Purification	bryo. At least two variants of a similar processed pseudo-					
Product:Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.Kestern blot analysis of extracts from HepG2,SW620 cells using BS48474 at 1: 1000.Molecular Weight:BS48474 at 1: 1000.29 kDaNote:Swiss-Prot:Note:095863For research use only, not for use in diagnostic procedurePurification&Purity:Affinity Purification	gene have been found on chromosome 2. [provided by					
Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.Molecular Weight:Western blot analysis of extracts from HepG2,SW620 cells using BS48474 at 1: 1000.Swiss-Prot:BS48474 at 1: 1000.O95863Note:Purification&Purity:For research use only, not for use in diagnostic procedured For research use only, not for use in diagnostic procedured For research use only, not for use in diagnostic procedured	RefSeq, Jul 2008]					
0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.Western blot analysis of extracts from HepG2,SW620 cells using BS48474 at 1: 1000.29 kDaBS48474 at 1: 1000.Swiss-Prot:Note:O95863For research use only, not for use in diagnostic procedurePurification&Purity:Affinity Purification	Product:					
0.05% BSA. Stable for 12 months from date of receipt.Western blot analysis of extracts from HepG2,SW620 cells usingMolecular Weight:Western blot analysis of extracts from HepG2,SW620 cells using29 kDaBS48474 at 1: 1000.Swiss-Prot:Note:O95863For research use only, not for use in diagnostic proceduredPurification&Purity:HepG2,SW620 cells usingAffinity PurificationHepG2,SW620 cells using	Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4),					
Molecular Weight: Western blot analysis of extracts from HepG2,SW620 cells using 29 kDa BS48474 at 1: 1000. Swiss-Prot: Note: O95863 For research use only, not for use in diagnostic procedure Purification&Purity: Affinity Purification	0.15M NaCl, 40%Glycerol, 0.01% sodium azide and					
29 kDaBS48474 at 1: 1000.Swiss-Prot:Note:095863For research use only, not for use in diagnostic proceduredPurification&Purity:For research use only, not for use in diagnostic proceduredAffinity PurificationFor research use only, not for use in diagnostic procedured	0.05% BSA. Stable for 12 months from date of receipt.					
Swiss-Prot: Note: O95863 For research use only, not for use in diagnostic procedure Purification&Purity: Affinity Purification	Molecular Weight:				Western blot analysis of extra	acts from HepG2,SW620 cells using
O95863 For research use only, not for use in diagnostic procedure Purification&Purity: Affinity Purification	29 kDa				BS48474 at 1: 1000.	
Purification&Purity: Affinity Purification	Swiss-Prot:					
Affinity Purification	O95863				For research use only,	not for use in diagnostic procedure.
•	Purification&Purity:					
	Affinity Purification					
Applications:	Applications:					

Biowo	rld Technology, Inc.			
Add:	1660 South Highway 100, Suite 500 St. Louis Park,			
	MN 55416,USA.			
Email:	info@bioworlde.com			
Tel:	6123263284			
Fax:	6122933841			