

p27 Kip1 (phospho-T198) polyclonal antibody

Catalog: BS4851

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

Cell cycle progression is regulated by a series of cyclin-dependent kinases that consist of catalytic subunits, designated Cdk, and activating subunits, designated cyclins. Orderly progression through the cell cycle requires the activation and inactivation of different cyclin-Cdk at appropriate times. A series of proteins has been recently described that function as “mitotic inhibitors.” These include p21, the levels of which are elevated upon DNA damage in G1 in a p53-dependent manner, p16 and a more recently described p16 related inhibitor designated p15. A p21 related protein, p27, has been described as a negative regulator of G1 progression and has been speculated to function as a possible mediator of TGFβ-induced G1 arrest. p27 interacts strongly with D-type cyclins and Cdk4 in vitro and to a lesser extent with cyclin E and Cdk2.

Product:

1 mg/ml in Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2.

Molecular Weight:

~ 27 kDa

Swiss-Prot:

P46527

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

p-p27 Kip1 (T198) polyclonal antibody detects endogenous levels of p27 Kip1 protein only when phosphorylated at Thr198.

DATA:



Western blot (WB) analysis of p-p27 Kip1 (T198) polyclonal antibody at 1:500 dilution

Lane1: HEK293T cell lysate treated with EGF(0.1ng/mL, 30mins)

Lane2: Raw264.7 cell lysate treated with EGF(0.1ng/mL, 30mins)

Lane3: H9C2 cell lysate treated with EGF(0.1ng/mL, 30mins)

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151