

Human, Mouse, Rat

# **Anterior Gradient 2 Recombinant Rabbit mAb**

Catal	og:	BS48	3515
-------	-----	------	------

Host: Rabbit

## BackGround:

This gene encodes a member of the disulfide isomerase (PDI) family of endoplasmic reticulum (ER) proteins that catalyze protein folding and thiol-disulfide interchange reactions. The encoded protein has an N-terminal ER-signal sequence, a catalytically active thioredoxin domain, and a C-terminal ER-retention sequence. This protein plays a role in cell migration, cellular transformation and metastasis and is as a p53 inhibitor. As an ER-localized molecular chaperone, it plays a role in the folding, trafficking, and assembly of cysteine-rich transmembrane receptors and the cysteine-rich intestinal gyl-coprotein mucin. This gene has been implicated in inflammatory bowel disease and cancer progression. [provided by RefSeq, Mar 2017]

#### **Product:**

Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.

#### **Molecular Weight:**

18 kDa

**Swiss-Prot:** 

#### 095994

**Purification&Purity:** 

Reactivity:

Affinity Purification

Applications:

WB: 1:1000<br />IHC: 1:200<br />ICC/IF: 1:100<br />FC: 1:20


#### **Storage&Stability:**

Store at  $4 \,^{\circ}{\rm C}$  short term. Aliquot and store at  $-20 \,^{\circ}{\rm C}$  long term. Avoid freeze-thaw cycles.

**Isotype:** 

### IgG

**DATA:** 



Western blot analysis of extracts from HeLa cells at 1:1000.

#### Note:

For research use only, not for use in diagnostic procedure.