



HDAC6 Recombinant Rabbit mAb

Catalog: BS49243

Host: Rabbit

Reactivity: Mouse, Rat

BackGround:

Enables cytoskeletal protein binding activity; protein deacetylase activity; and ubiquitin binding activity. Involved in several processes, including positive regulation of peptidyl-serine phosphorylation; protein deacetylation; and protein destabilization. Acts upstream of or within several processes, including aggresome assembly; cellular protein metabolic process; and neuron projection morphogenesis. Located in several cellular components, including dendrite; microtubule cytoskeleton; and perikaryon. Part of protein-containing complex. Is expressed in several structures, including central nervous system; early embryo; future midbrain; gut gland; and lung. Human ortholog(s) of this gene implicated in chondrodysplasia with platyspondyly, distinctive brachydactyly, hydrocephaly, and microphthalmia. Orthologous to human HDAC6 (histone deacetylase 6). [provided by Alliance of Genome Resources, Apr 2022]

Product:

Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.

Molecular Weight:

160 kDa

Swiss-Prot:

Q9Z2V5

Purification&Purity:

Affinity Purification

Applications:

WB: 1:1000
FC: 1:500
IP: 1:20


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Isotype:

IgG

DATA:



Western blot analysis of extracts from Neuro-2a cells at 1:1000.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park,
MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046,
P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151