

# **USP19** polyclonal antibody

Catalog: BS5975

Host: Rabbit

Reactivity: Human, Mouse, Rat

# **BackGround:**

The ubiquitin (Ub) pathway involves three sequential enzymatic steps that facilitate the conjugation of Ub and Ub-like molecules to specific protein substrates. Through the use of a wide range of enzymes that can add or remove ubiquitin, the Ub pathway controls many intracellular processes such as signal transduction, transcriptional activation and cell cycle progression. Ubiquitin-specific-processing protease 19 (USP19), also known as ubiquitin carboxyl-terminal hydrolase 19, ubiquitin thioesterase 19 and deubiquitinating enzyme 19, is a 1318 amino acid member of the peptidase C19 family of proteins. USP19 is thought to be involved in the ubiquitin-dependent proteolytic pathway in conjunction with the 26S proteasome. Four known isoforms of USP19 exist as a result of alternative splicing events.

# **Product:**

1 mg/ml in Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2.

# **Molecular Weight:**

~ 146 kDa

**Swiss-Prot:** 

094966

#### **Purification&Purity:**

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

#### **Applications:**

WB: 1:500~1:1000

IHC: 1:50~1:200


Storage&Stability:

Store at  $4 \,^{\circ}{\rm C}$  short term. Aliquot and store at  $-20 \,^{\circ}{\rm C}$  long term. Avoid freeze-thaw cycles.

## **Specificity:**

USP19 polyclonal antibody detects endogenous levels of USP19 protein.


**DATA:** 



Western blot (WB) analysis of USP19 polyclonal antibody at 1:500 dilution

Lane1:MCF-7 cell lysate Lane2:HEK293T cell lysate

Lane3:H9C2 cell lysate



Immunohistochemistry (IHC) analyzes of USP19 pAb in paraffin-embedded human breast carcinoma tissue at 1:50.showing endoplasmic reticulum membrane staining. Negative control (the right)Using PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin followed by avidin-peroxidase.

#### Note:

For research use only, not for use in diagnostic procedure.

# Bioworld Technology, Inc.

 
 Add:
 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA.

 Email:
 info@bioworlde.com

 Tel:
 6123263284

 Fax:
 6122933841

# Bioworld technology, co. Ltd.

 
 Add:
 No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

 Email:
 info@biogot.com

 Tel:
 0086-025-68037686

 Fax:
 0086-025-68035151