

BCAT1 polyclonal antibody

Catalog: BS60374

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

ECA39, also known as BCAT1 (branched-chain-amino-acid aminotransferase 1, cytosolic) is localized to the cytoplasm where it forms a homodimer. ECA39 is expressed in the brain and kidney during embryogenesis and is overexpressed in c-Myc induced tumors. BCAT2 (branched-chain-amino-acid aminotransferase 2, mitochondrial), also known as placental protein 18 (PP18), is expressed as two isoforms produced by alternative splicing. The first isoform of BCAT2, designated BCAT2A, is expressed in the mitochondrion, while the second isoform, designated BCAT2B, is expressed in the cytoplasm. Ubiquitously expressed, BCAT2 is also thought to act as a transporter of branched chain alpha-keto acids.

Product:

1mg/ml in PBS with 0.1% Sodium Azide, 50% Glycerol.

Molecular Weight:

~ 43 kDa

Swiss-Prot:

P54687

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

BCAT1 polyclonal antibody detects endogenous levels of BCAT1 protein.

DATA:



Western blot (WB) analysis of BCAT1 polyclonal antibody at 1:500 dilution

Lane1:Hela cell lysate

Lane2:NIH-3T3 cell lysate

Lane3:PC12 cell lysate

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogol.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151