

TBK1 polyclonal antibody

Catalog: BS60714

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

The transcription factor NF κ B is retained in the cytoplasm in an inactive form by the inhibitory protein I κ B. Activation of NF κ B requires that I κ B be phosphorylated on specific serine residues, which results in targeted degradation of I κ B. I κ B kinase α (IKK α), previously designated CHUK, interacts with I κ B- α and specifically phosphorylates I κ B- α on the sites that trigger its degradation, serines 32 and 36. The functional IKK complex contains three subunits, IKK α , IKK β and IKK γ (also designated NEMO), and each appear to make essential contributions to I κ B phosphorylation. TANK binding kinase (TBK1), also designated T2K, is a novel IKK-related kinase that has been identified in murine and human tissues. TBK1 was shown to complex with TRAF2 and TANK in the NF κ B activation pathway. TBK1 shares homology with IKK α and IKK β in the amino-terminal half, which includes the kinase domain.

Product:

1 mg/ml in Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.2.

Molecular Weight:

~ 84 kDa

Swiss-Prot:

Q9UHD2

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific im-

munogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

TBK1 polyclonal antibody detects endogenous levels of TBK1 protein.

DATA:



Western blot (WB) analysis of TBK1 polyclonal antibody at 1:500 dilution

Lane1:SK-OVCAR3 whole cell lysate(40ug)

Lane2:HCT116 whole cell lysate(40ug)

Lane3:CT26 whole cell lysate(40ug)

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151