

PLC γ 2 polyclonal antibody

Catalog: BS61288

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

Phosphoinositide-specific phospholipase C (PLC) plays a critical role in the initiation of receptor mediated signal transduction through the generation of the two second messengers, inositol 1,4,5-triphosphate and diacylglycerol from phosphatidylinositol 4,5-bisphosphate. There are many mammalian PLC isozymes, including PLC β 1, PLC β 2, PLC β 3, PLC β 4, PLC γ 1, PLC γ 2, PLC δ 1, PLC δ 2 and PLC ϵ . After stimulation of the collagen receptor glycoprotein VI in human platelets, PLC γ 2 associates with several tyrosine-phosphorylated proteins (Syk, SLP-76, Lyn, linker for activation of T cells (LAT) and the FcR γ chain), which bind to its C-terminal SH2 domain. PLC γ 1 associates with Syk in B cells, but PLC γ 2 does not associate with Syk in platelets. The C-terminal SH2 domain is involved in the regulation of PLC γ 2. In addition, Btk can induce PLC γ 2 tyrosine phosphorylation and initiate calcium mobilization in CD72-stimulated B lymphocytes.

Product:

1 mg/ml in Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.3.

Molecular Weight:

~ 148 kDa

Swiss-Prot:

P16885

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

PLC γ 2 polyclonal antibody detects endogenous levels of PLC γ 2 protein.

DATA:



Western blot (WB) analysis of PLC γ 2 polyclonal antibody at 1:500

Line1:HEK293T whole cell lysate

Line2:H9C2 whole cell lysate

Line3:Raw264.7 whole cell lysate

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151