

BS62498

Host:

Rabbit

Catalog:

Reactivity: Human, Mouse, Rat

GTF3A polyclonal antibody

Culuiog	100001110j
BackGround:	term. Avoid freeze-thaw cycles.
Transcription factor IIIA, involved in ribosomal large	Specificity:
subunit biogenesis. Binds the approximately 50 base pairs	GTF3A polyclonal antibody detects endogenous levels of
internal control region (ICR) of 5S ribosomal RNA genes.	GTF3A protein.
It is required for their RNA polymerase III-dependent	DATA:
transcription and may also maintain the transcription of	kDa A B C D
other genes. Also binds the transcribed 5S RNA's.	100
Product:	70
Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide,	50 35
50% glycerol, pH7.2	25
Molecular Weight:	20
~ 37 kDa	15
Swiss-Prot:	Western blot (WB) analysis of GTF3A polyclonal antibody at 1:500 di-
Q92664	lution
Purification&Purity:	LaneA:A549 whole cell lysate
The antibody was affinity-purified from rabbit antiserum	LaneB:COS-7 whole cell lysate
by affinity-chromatography using epitope-specific im-	LaneC:PC12 whole cell lysate
munogen and the purity is $> 95\%$ (by SDS-PAGE).	LaneD:AML-12 whole cell lysate
	Madaa

Applications:

WB:1:500~1:1000

Storage&Stability:

Store at 4 $^{\rm C}$ short term. Aliquot and store at -20 $^{\rm C}$ long

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.	
Add:	1660 South Highway 100, Suite 500 St. Louis Park,
	MN 55416,USA.
Email:	info@bioworlde.com
Tel:	6123263284
Fax:	6122933841