

MAX (phospho-S11) polyclonal antibody

Catalog: BS64009

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

It is now well established that the nature and relative abundance of individual subunits of different classes of transcription factors can positively or negatively regulate levels of gene expression. Myc proteins homodimerize and bind DNA poorly, if at all, at physiological levels. Max is a nuclear localized bHLH-Zip protein initially identified by screening a B cell expression library with the bHLH-Zip region of c-Myc. Max homodimers and the Myc-Max heterodimers bind the sequence CACGTG; however the binding of the hetero-dimeric complex is stronger than the Max homodimer. The Max gene products have been identified as Max and Max 9 proteins that differ by a 9 amino acid insertion N-terminal to the basic region. In contrast to Myc, which is highly regulated during progression through the cell cycle, Max is highly stable and is much more abundant than Myc. Two members of the bHLH-Zip protein family, designated Mad and Mxi1, homodimerize poorly but form heterodimeric complexes with Max that have opposing functions to Myc-Max heterodimers with respect to regulation of gene expression.

Product:

1mg/ml in PBS with 0.1% Sodium Azide, 50% Glycerol.

Molecular Weight:

~ 18 kDa

Swiss-Prot:

P61244

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific im-

munogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

p-MAX(S11) polyclonal antibody detects endogenous levels of MAX protein only when phosphorylated at Ser11.

DATA:



Western blot (WB) analysis of p-MAX (S11) polyclonal antibody at 1:500 dilution

Lane1:Hela whole cell lysate

Lane2:Mouse heart tissue lysate

Lane3:H9C2 whole cell lysate

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151