

Bax polyclonal antibody

Catalog: BS6420

Host:

Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

The Bcl-2 gene was isolated at the chromosomal breakpoint of t-bearing follicular B cell lymphomas. Bcl-2 blocks cell death following a variety of stimuli and confers a death-sparing effect to certain hematopoietic cell lines following growth factor withdrawal. Bcl-2 is localized to outer mitochondrial membranes and endoplasmic reticulum as well as nuclear membranes. A related protein, designated Bax p21 (for Bcl-associated X protein), has extensive amino acid homology with Bcl-2 and both homodimerizes and forms heterodimers with Bcl-2. Overexpression of Bax accelerates apoptotic death induced by cytokine deprivation in an IL-3 dependent cell line, and Bax also counters the death repressor activity of Bcl-2.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 20 kDa

Swiss-Prot:

Q07812

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB 1:500 - 1:1000
IHC 1:50 - 1:200

Storage&Stability:

Store at $4 \,^{\circ}{\rm C}$ short term. Aliquot and store at $-20 \,^{\circ}{\rm C}$ long term. Avoid freeze-thaw cycles.

Specificity:

Bax polyclonal antibody detects endogenous levels of Bax protein.

DATA:


Bioworld Technology, Inc.

 Add:
 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA.

 Email:
 info@bioworlde.com

 Tel:
 6123263284

 Fax:
 6122933841




Western blot analysis of extracts of various cell lines, using BAX anti-

body at 1:1000 dilution.

Lysates/proteins: 40ug per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Exposure time: 90s.




Western blot analysis of extracts of various cell lines, using BAX antibody at 1:1000 dilution.

Lysates/proteins: 40ug per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Exposure time: 90s.




Immunohistochemistry of paraffin-embedded mouse testis using BAX antibody at dilution of 1:100 (40x lens).

Bioworld technology, co. Ltd. Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China. Email: info@biogot.com Tel: 0086-025-68037686 Fax: 0086-025-68035151



PRODUCT DATA SHEET

Bioworld Technology,Inc.



Immunoprecipitation - [KO Validated] Bax Rabbit pAb

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.		Bioworld technology, co. Ltd.	
Add:	1660 South Highway 100, Suite 500 St. Louis Park,	Add:	No 9, weidi road Qixia District Nanjing, 210046,
	MN 55416,USA.		P. R. China.
Email:	info@bioworlde.com	Email:	info@biogot.com
Tel:	6123263284	Tel:	0086-025-68037686
Fax:	6122933841	Fax:	0086-025-68035151