

CREB-1 Polyclonal Antibody

Catalog: BS65863

Host: Rabbit

Reactivity: Human, Mouse, Rat,

BackGround:

Cyclic AMP Response Element Binding protein (CREB) is a basic / leucine zipper transcription factor that binds the cyclic AMP response element (CRE) and activates transcription in response to a variety of extracellular signals including neurotransmitters, hormones, membrane depolarization, and growth and neurotrophic factors. Activation of CREB is dependent upon the phosphorylation of serine 133. Phosphorylation occurs via p44 / 42 MAP kinase and p90RSK and also via p38 MAP kinase and MSK 1. Although CREB will bind DNA independent of its phosphorylation state, only the phosphorylated form is competent as a transcription factor. CREB binding protein (CBP), a transcriptional coactivator that directly interacts with CREB, binds to CREB in the region of serine 133. CREB appears to play an important role in learning and memory. CREB knock out mice show diminished learning ability.

Product:

0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

Molecular Weight:

~47 kDa

Swiss-Prot:

P16220

Purification&Purity:

affinity purified by Protein A

Applications:

WB=1:500-2000


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -36 °C long term. Avoid freeze-thaw cycles.


Specificity:

CREB-1 Polyclonal Antibody detects endogenous levels of CREB-1 protein.

DATA:



Primary: Anti-CREB-1 at 1/300 dilution



Primary: Anti-CREB-1 at 1/1000 dilution

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogol.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151