

IL13 polyclonal antibody

Catalog: BS6637

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

Interleukin-13, or IL-13, is a pleiotropic cytokine that exhibits 30% sequence identity with IL-4 and shares many of the same biological activities. Like IL-4, IL-13 affects monocytes, macrophages and B cells by upregulating the expression of CD23 and MHC proteins, and downregulating the expression of CD14. Both IL-4 and IL-13 are secreted by activated T lymphocytes and are powerful regulators of inflammation. Both inhibit the secretion of proinflammatory cytokines and chemokines from activated monocytes and stimulate the expression of IgE on activated B cells. IL-13 contains five Cysteine residues and multiple N-linked glycosylation sites and has been reported to inhibit the production of IL-2 in natural killer cells. IL-13 cDNA encodes a 131 amino acid precursor with a 20 amino acid signal peptide which is cleaved to generate a mature protein.

Product:

1mg/ml in PBS with 0.1% Sodium Azide, 50% Glycerol.

Molecular Weight:

~ 16 kDa

Swiss-Prot:

P35225

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific im-

munogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:2000

IHC: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

IL13 polyclonal antibody detects endogenous levels of IL13 protein.

DATA:



Western blot analysis of extracts of recombinant protein using IL13 antibody.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151