

## PRODUCT DATA SHEET

Bioworld Technology,Inc.

# Caspase-3 p17 (D175) polyclonal antibody

Catalog: BS7073 Host: Rabbit Reactivity: Human, Mouse, Rat

#### **BackGround:**

Caspase 3 (also known as CPP32, YAMA and apopain) is the most extensively studied apoptotic protein among caspase family members. Caspase 3 is synthesized as inactive pro enzyme that is processed in cells undergoing apoptosis by self proteolysis and/or cleavage by other upstream proteases (e.g. Caspases 8, 9 and 10). The processed form of Caspase 3 consists of large (17kD) and small (12kD) subunits which associate to form an active enzyme. Caspase 3 is cleaved at Asp28 - Ser29 and Asp175 - Ser176. The active Caspase 3 proteolytically cleaves and activates other caspases (e.g. Caspases 6, 7 and 9), as well as relevant targets in the cells (e.g. PARP and DFF).

## **Product:**

1mg/ml in PBS with 0.1% Sodium Azide, 50% Glycerol.

#### **Molecular Weight:**

~ 17 kDa

#### **Swiss-Prot:**

P42574

## **Purification&Purity:**

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

#### **Applications:**

WB: 1:500~1:1000

#### **Storage&Stability:**


Store at  $4 \,\mathrm{C}$  short term. Aliquot and store at  $-20 \,\mathrm{C}$  long term. Avoid freeze-thaw cycles.

## **Specificity:**

Caspase-3 p17 (D175) polyclonal antibody detects endogenous levels of the large fragment (17 kDa) of activated caspase-3 resulting from cleavage adjacent to Asp175. This antibody does not recognize full length

caspase-3 or other cleaved caspases.

## **DATA:**




Western blot (WB) analysis of Caspase-3 p17 (D175) polyclonal anti-

body at 1:500 dilution

Lane1:Hela cell lysate treated with Lps

Lane2:Raw264.7 cell lysate treated with Lps

Lane3:PC12 cell lysate treated with Lps



Western blot (WB) analysis of Caspase-3 p17 (D175) pAb at 1:500 dilution

Lane1:HepG2 whole cell lysate(40ug)

Lane2:HepG2 treated with Etoposide (20ng/ml, 20h) whole cell lysate (40ug)

Lane3:HepG2 treated with Etoposide (20ng/ml, 20h, $\lambda$ -ppase) whole cell lysate (40ug)

#### Note:

For research use only, not for use in diagnostic procedure.

## Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park,

MN 55416,USA.

Email: <u>info@bioworlde.com</u>

Tel: 6123263284 Fax: 6122933841 Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046,

P. R. China.

Email: <u>info@biogot.com</u> Tel: 0086-025-68037686 Fax: 0086-025-68035151