

USP28 polyclonal antibody

Catalog: BS71629 Host: Rabbit	Reactivity: Human, Mouse
BackGround:	WB 1:500 - 1:2000
The ubiquitin-dependent protein degradation pathway is	Storage&Stability:
essential for proteolysis of intracellular proteins and pep-	Store at 4 ${\rm C}$ short term. Aliquot and store at -20 ${\rm C}$ long
tides. Enzymes that remove ubiquitin from ubiqui-	term. Avoid freeze-thaw cycles.
tin-conjugated peptides, like USP28, affect the fate and	Specificity:
degradation of intracellular proteins and are essential for	USP28 polyclonal antibody detects endogenous levels of
maintenance of cell-free ubiquitin pools (Valero et al.,	USP28 protein.
2001).	DATA:
Product:	rrain

1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Molecular Weight:

Predicted band size: 118 kDa

Observed band size: Refer to Figures


Swiss-Prot:

Q96RU2

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:



Western blot analysis of extracts of various cell lines, using USP28 an-

tibody.

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.	
Add:	1660 South Highway 100, Suite 500 St. Louis Park,
	MN 55416,USA.
Email:	info@bioworlde.com
Tel:	6123263284
Fax:	6122933841