

Smad4 polyclonal antibody

Catalog: BS7225

Host: Rabbit

Reactivity:

ity: Human, Mouse

BackGround:

Smad proteins, the mammalian homologs of the Drosophila Mothers against dpp (Mad) have been implicated as downstream effectors of TGF β /BMP signaling. Smad1 (also designated Madr1 or JV4-1), Smad5 and mammalian Smad8 (also designated Smad9 or MadH6) are effectors of BMP2 and BMP4 function while Smad2 (also designated Madr2 or JV18-1) and Smad3 are involved in TGF β and activin-mediated growth modulation. Smad4 (also designated DPC4) has been shown to mediate all of the above activities through interaction with various Smad family members. Smad6 and Smad7 regulate the response to activin/TGF β signaling by interfering with TGF β -mediated phosphorylation of other Smad family members.

Product:

1mg/ml in PBS with 0.1% Sodium Azide, 50% Glycerol. Molecular Weight:

~ 60 kDa

Swiss-Prot:

Q13485

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:2000 IF: 1:50~1:200

Storage&Stability:

Store at $4 \,^{\circ}{\rm C}$ short term. Aliquot and store at -20 $^{\circ}{\rm C}$ long term. Avoid freeze-thaw cycles.


Specificity:

Smad4 polyclonal antibody detects endogenous levels of Smad4 protein.

DATA:



Western blot analysis of Smad4 polyclonal antibody



Immunofluorescence analysis of U2OS cells, using Smad4 polyclonal antibody

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

 Add:
 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA.

 Email:
 info@bioworlde.com

 Tel:
 6123263284

 Fax:
 6122933841

Bioworld technology, co. Ltd.

 Add:
 No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

 Email:
 info@biogot.com

 Tel:
 0086-025-68037686

 Fax:
 0086-025-68035151