

#### PRODUCT DATA SHEET

Bioworld Technology,Inc.

# WNT10B polyclonal antibody

Catalog: BS76917 Host: Rabbit Reactivity: Human, Mouse, Rat

#### **BackGround:**

The WNT gene family consists of structurally related genes which encode secreted signaling proteins. These proteins have been implicated in oncogenesis and in several developmental processes, including regulation of cell fate and patterning during embryogenesis. This gene is a member of the WNT gene family. It may be involved in breast cancer, and its protein signaling is likely a molecular switch that governs adipogenesis. This protein is 96% identical to the mouse Wnt10b protein at the amino acid level. This gene is clustered with another family member, WNT1, in the chromosome 12q13 region.

#### **Product:**

1 mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

#### **Molecular Weight:**

43kDa

#### **Swiss-Prot:**

O00744

## **Purification&Purity:**

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

#### **Applications:**

WB,1:500 - 1:2000|IHC,1:50 - 1:200|IF/ICC,1:50 - 1:200


#### Storage&Stability:

Store at  $4\,\mathrm{C}$  short term. Aliquot and store at  $-20\,\mathrm{C}$  long term. Avoid freeze-thaw cycles.


#### **Modification:**

Unmodification


#### **DATA:**



Western blot analysis of extracts of HeLa cells, using WNT10B antibody at 1:1000 dilution.<br/>
Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.<br/>
br/>Lysates/proteins: 25ug per lane.<br/>
Blocking buffer: 3% nonfat dry milk in TBST.<br/>
Detection: ECL Basic Kit .<br/>
Exposure time: 10s.



Immunohistochemistry of paraffin-embedded rat ovary using WNT10B antibody at dilution of 1:100 .Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry of paraffin-embedded human colon carcinoma using WNT10B antibody at dilution of 1:100 .Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.

# Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park,

MN 55416,USA.

Email: <u>info@bioworlde.com</u>

Tel: 6123263284 Fax: 6122933841

# Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046,


P. R. China.

Email: <u>info@biogot.com</u>
Tel: 0086-025-68037686
Fax: 0086-025-68035151



# PRODUCT DATA SHEET

Bioworld Technology,Inc.



Immunofluorescence analysis of C6 cells using WNT10B antibody at dilution of 1:100. Blue: DAPI for nuclear staining.

## Note:

For research use only, not for use in diagnostic procedure.

# Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park,

MN 55416,USA.

Email: <u>info@bioworlde.com</u>

Tel: 6123263284 Fax: 6122933841

# Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046,

P. R. China.

Email: <u>info@biogot.com</u>
Tel: 0086-025-68037686
Fax: 0086-025-68035151