

## SLC8A1 polyclonal antibody

Catalog: BS78863

Host: Rabbit

Reactivity: Human

### BackGround:

In cardiac myocytes,  $\text{Ca}^{2+}$  concentrations alternate between high levels during contraction and low levels during relaxation. The increase in  $\text{Ca}^{2+}$  concentration during contraction is primarily due to release of  $\text{Ca}^{2+}$  from intracellular stores. However, some  $\text{Ca}^{2+}$  also enters the cell through the sarcolemma (plasma membrane). During relaxation,  $\text{Ca}^{2+}$  is sequestered within the intracellular stores. To prevent overloading of intracellular stores, the  $\text{Ca}^{2+}$  that entered across the sarcolemma must be extruded from the cell. The  $\text{Na}^{+}$ - $\text{Ca}^{2+}$  exchanger is the primary mechanism by which the  $\text{Ca}^{2+}$  is extruded from the cell during relaxation. In the heart, the exchanger may play a key role in digitalis action. The exchanger is the dominant mechanism in returning the cardiac myocyte to its resting state following excitation.

### Product:

1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

### Molecular Weight:

109kDa

### Swiss-Prot:

P32418

### Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific im-

munogen and the purity is > 95% (by SDS-PAGE).

### Applications:

WB, 1:500 - 1:2000


### Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

### Modification:

Unmodification

### DATA:



Western blot analysis of extracts of U-87MG cells, using SLC8A1 antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 1s.

### Note:

For research use only, not for use in diagnostic procedure.

### Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: [info@bioworld.com](mailto:info@bioworld.com)

Tel: 6123263284

Fax: 6122933841

### Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: [info@biogot.com](mailto:info@biogot.com)

Tel: 0086-025-68037686

Fax: 0086-025-68035151