

ZCCHC8 polyclonal antibody

Catalog: BS79242

Host: Rabbit

Reactivity: Human, Mouse

BackGround:

This gene encodes a scaffold protein which serves as an accessory factor to the nuclear RNA exosome complex. The encoded protein forms a trimeric human nuclear exosome targeting (NEXT) complex, together with hMTR4 and the RNA-binding factor RBM7 which promotes the exosomal degradation of non-coding promoter-upstream transcripts, enhancer RNAs and 3'-extended products of histone- and small nuclear RNA transcription. This complex is also thought to recruit the exosome to degrade intronic RNAs via its interaction with both the exosome and the spliceosome. It contains both an N-terminal zinc-knuckle domain and a C-terminal proline-rich domain.

Product:

1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

78kDa

Swiss-Prot:

Q6NZY4

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:


WB, 1:500 - 1:2000

Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Modification:

Unmodification

DATA:

Western blot analysis of extracts of various cell lines, using ZCCHC8 antibody at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 15s.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151