

## Pan-Succinyl-K polyclonal antibody

Catalog: BS79342

Host: Rabbit

Reactivity: Human, Mouse, Rat

### Background:

Succinylation of protein is a kind of post-translational modification of protein mainly occurring on lysine residues. Compared with methylation and acetylation, succinylation of lysine has a greater effect on protein structure and function. Sirt5 is an important de-succinylation modifying enzyme involved in the regulation of a variety of biological processes, such as reactive oxygen species defense, fatty acid metabolism and cell apoptosis

### Product:

1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

### Molecular Weight:

10-150KDa

### Swiss-Prot:

### Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

### Applications:

WB, 1:500 - 1:2000


### Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

### Modification:

Other Modified

### DATA:



Western blot analysis of extracts of various cell lines, using Pan-Succinyl-K antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 180s.

### Note:

For research use only, not for use in diagnostic procedure.

### Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: [info@bioworld.com](mailto:info@bioworld.com)

Tel: 6123263284

Fax: 6122933841

### Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: [info@biogot.com](mailto:info@biogot.com)

Tel: 0086-025-68037686

Fax: 0086-025-68035151