

149kDa

PLC gamma 1 (PLCG1) polyclonal antibody

Catalog:	BS79977	Host:	Rabbit	Reactivity: Human, Mouse, Rat
BackGround:				Swiss-Prot:
The protein encoded by this gene catalyzes the formation				P19174
of inositol 1,4,5-trisphosphate and diacylglycerol from				Purification&Purity:
phosphatidylinositol 4,5-bisphosphate. This reaction uses			reaction uses	The antibody was affinity-purified from rabbit antiserum
calcium as a cofactor and plays an important role in the			nt role in the	by affinity-chromatography using epitope-specific im-
intracellular transduction of receptor-mediated tyrosine			ated tyrosine	munogen and the purity is $> 95\%$ (by SDS-PAGE).
kinase activators. For example, when activated by SRC,			ted by SRC,	Applications:
the encoded protein causes the Ras guanine nucleotide exchange factor RasGRP1 to translocate to the Golgi, where it activates Ras. Also, this protein has been shown to be a major substrate for heparin-binding growth factor 1 (acidic fibroblast growth factor)-activated tyrosine ki- nase. Two transcript variants encoding different isoforms have been found for this gene.			e nucleotide	WB,1:500 - 1:2000
			o the Golgi,	Storage&Stability:
			been shown	Store at 4 ${}^\circ\!\!\!{\rm C}$ short term. Aliquot and store at -20 ${}^\circ\!\!\!{\rm C}$ long
			growth factor	term. Avoid freeze-thaw cycles.
			•	Modification:
			ent isoforms	Unmodification
				DATA:
Product:				
1mg/ml in PBS with 0.02% sodium azide, 50% glycerol,			0% glycerol,	Note:
pH7.2				For research use only, not for use in diagnostic procedure.
Molecular Weight:				