

elF4E polyclonal antibody

Catalog:	BS80464	Host:	Rabbit	Reactivity:	Human, Mouse, Rat
BackGround: The protein encoded by this gene is a component of the eukaryotic translation initiation factor 4F complex, which recognizes the 7-methylguanosine cap structure at the 5' end of messenger RNAs. The encoded protein aids in translation initiation by recruiting ribosomes to the 5'-cap structure. Association of this protein with the 4F complex is the rate-limiting step in translation initiation. This gene acts as a proto-oncogene, and its expression and activation is associated with transformation and tumorigenesis.				munogen and the purity Applications: WB,1:500 - 1:2000 IF/I Storage&Stability:	y is > 95% (by SDS-PAGE). CCC,1:50 - 1:200 h. Aliquot and store at -20 ℃ long
Several pseudogenes of this gene are found on other chromosomes. Alternative splicing results in multiple transcript variants. Product:			in multiple	Western blot analysis of extracts of various cell lines, using eIF4E anti- body at 1:1000 dilution. br/>Secondary antibody: HRP Goat An- ti-Rabbit IgG at 1:10000 dilution. Lysates/proteins: 25ug per 	
pH7.2	ng/ml in PBS with 0.02% sodium azide, 50% glycerol, H7.2 Iolecular Weight:				
Molecular					
25kDa				Note:	
Swiss-Prot:				For research use only, not for use in diagnostic procedure.	
P06730					

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific im-

Bioworld Technology, Inc. Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA. Email: info@bioworlde.com Tel: 6123263284 Fax: 6122933841

Bioworld technology, co. Ltd. Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China. Email: <u>info@biogot.com</u> Tel: 0086-025-68037686 Fax: 0086-025-68035151