

NNMT polyclonal antibody

Catalog: BS8344

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

Nicotinamide N-methyltransferase (NNMT) catalyzes the N-methylation of nicotin-amide and other pyridines. NNMT activity in the human liver has a bimodal frequency distribution, indicating that its enzyme activity may be modulated through a genetic polymorphism, which could have functional implications for individual differences in drug and xenobiotic toxicity. The gene that encodes human NNMT is approximately 16.5 kb in length, consists of three exons and two introns and maps to 11q23.2. NNMT isolated from the human liver was determined to be 969 nucleotides in length, with a 792 nucleotide open reading frame that encodes a 264 amino acid protein. The NNMT gene is presumed to be a significant genetic determinant of plasma homocysteine levels in Spanish families, since it encodes an enzyme involved in homocysteine synthesis.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 30 kDa

Swiss-Prot:

P40261

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:2000

IF: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

NNMT polyclonal antibody detects endogenous levels of NNMT protein.

DATA:



WesternBlot (WB) analysis of NNMT polyclonal antibody

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151