

PI3K p110 γ (L89) polyclonal antibody

Catalog: BS9175

Host: Rabbit

Reactivity: Human

BackGround:

Phosphatidylinositol 3-kinase (PI 3-kinase) is composed of p85 and p110 subunits. p85 lacks PI 3-kinase activity and acts as an adapter, coupling p110 to activated protein tyrosine kinase. Two forms of p85 have been described (p85 α and p85 β), each possessing one SH3 and two SH2 domains. Various p110 isoforms have been identified. p110 α and p110 β interact with p85 α , and p110 α has also been shown to interact with p85 β in vitro. p110 γ expression is restricted to white blood cells. It has been shown to bind p85 α and β , but it apparently does not phosphorylate these subunits. p110 γ seems to have the capacity to autophosphorylate. p110 does not interact with the p85 subunits. It has been shown to be activated by α and β heterotrimeric G proteins.

Product:

1 mg/ml in Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2.

Molecular Weight:

~ 110 kDa

Swiss-Prot:

P48736

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:


PI 3-Kinase p110 γ (L89) polyclonal antibody detects endogenous levels of PI 3-Kinase p110 γ protein.

DATA:




BS9175
Lot CA36131

Immunohistochemistry (IHC) analyzes of PI3K p110 γ (L89) pAb in paraffin-embedded human breast carcinoma tissue at 1:50, showing strong cytoplasmic staining. Negative control (the right) Using PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin followed by avidin-peroxidase.



BS9175
Lot CA36131

Immunohistochemistry (IHC) analyzes of PI3K p110 γ (L89) pAb in paraffin-embedded human lung carcinoma tissue at 1:50, showing strong cytoplasmic staining. Negative control (the right) Using PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin followed by avidin-peroxidase.



BS9175
Lot CA36131

Immunohistochemistry (IHC) analyzes of PI3K p110 γ (L89) pAb in paraffin-embedded human kidney carcinoma tissue at 1:50, showing strong cytoplasmic staining. Negative control (the right) Using PBS in-

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151



PRODUCT DATA SHEET

Bioworld Technology, Inc.

stead of primary antibody, secondary antibody is Goat Anti-Rabbit
IgG-biotin followed by avidin-peroxidase.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park,
MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046,
P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151