

CD9 polyclonal antibody

BackGround:	
Involved in platelet activation and aggregation. Regulates paranodal junction formation. Involved in cell adhesion, cell motility and tumor metastasis. Required for sperm-egg fusion. Product: 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol,	kDa JST-M ^C 250- 150- 100- 75- 50- 37- 25- 20- 15- 10- 10- 10- 10- 10- 10- 15- 10- 10- 10- 10- 10- 10- 10- 10
0.01% Sodium azide and 0.05% BSA Molecular Weight: Calculated MW: 25 kDa; Observed MW: 25 kDa Swiss-Prot: P21926 Purification&Purity: Affinity Purified Applications: WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 IP: 1/20 FC: 1/50-1/100 Storage&Stability: Store at 4 °C short term. Aliquot and store at -20 °C long	Western blot analysis of CD9 in U87-MG lysates using CD9 antibody. Image: CD9 in U87-MG lysates using CD9 antibody. Immunocytochemistry analysis of CD9 in Hela using CD9 antibody, and DAPI Immunohistochemistry analysis of paraffin-embedded Human Cholangiocarcinoma using CD9 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.
term. Avoid freeze-thaw cycles.	Note:
Isotype: IgG DATA:	For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

1660 South Highway 100, Suite 500 St. Louis Park,
MN 55416,USA.
info@bioworlde.com
6123263284
6122933841

Bioworld technology, co. Ltd. Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China. Email: <u>info@biogot.com</u> Tel: 0086-025-68037686 Fax: 0086-025-68035151