

PRODUCT DATA SHEET

Bioworld Technology,Inc.

PER2 polyclonal antibody

Catalog: GCP101 Host: Rabbit Reactivity: Human, Pig

BackGround:

Biological timepieces called circadian Clocks are responsible for the regulation of hormonal rhythms, sleep cycles and other behaviors. The superchiasmaticnucleus (SCN), which is located in the brain, was the first mammalian circadian Clock to be discovered. A number of transcription factors appearing to bemolecular components of the SCN Clock have been identified. Mutations within the Clock gene increase the length of the endogenous period and cause aloss of rhythmicity of circadian oscillations. Three mammalian period proteins, designated Per1, Per2 and Per3, exhibit circadian rhythyms in the SCN. Duringsubjective night, Per1 and Per2 RNA levels increase in response to light pulses while Per3 RNA levels show no change in response to light pulses. Tim, fortimeless, interacts with Per1 as well as Per2; and Tim and Per1 negativelyregulate Clock-BMAL1-induced transcription.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2.

Molecular Weight:

~ 155 kDa

Swiss-Prot:

O15055

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:2000~1:5000 IHC: 1:50~1:200

Storage&Stability:


Store at $4\,\mathrm{C}$ short term. Aliquot and store at $-20\,\mathrm{C}$ long term. Avoid freeze-thaw cycles.

Specificity:

PER2 polyclonal antibody detects endogenous levels of

PER2 protein.

DATA:



Western blot (WB) analysis of PER2 polyclonal antibody at 1:2000 di-

lution


Lane1:A549 whole cell lysate(40ug)

Lane2:The Spleen tissue lysate of Pig(40ug)


Lane3:The Lung tissue lysate of Pig(40ug)

Lane4:MCF-7 whole cell lysate(40ug)

Lane5:Hela whole cell lysate(40ug)



Immunohistochemistry of paraffin-embedded Human Brain using PER2 antibody at dilution of 1:50.



Western blot (WB) analysis of PER2 pAb at 1:1000 dilution

Lane1:A549 whole cell lysate(40ug)

Lane2:PER2 knockdown A549 whole cell lysate(40ug)

Note:

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park,

MN 55416,USA.

Email: info@bioworlde.com

Tel: 6123263284 Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046,

P. R. China.

Email: info@biogot.com
Tel: 0086-025-68037686
Fax: 0086-025-68035151



PRODUCT DATA SHEET

Bioworld Technology,Inc.

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park,

MN 55416,USA.

Email: info@bioworlde.com

Tel: 6123263284 Fax: 6122933841 Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046,

P. R. China.

Email: info@biogot.com
Tel: 0086-025-68037686
Fax: 0086-025-68035151