

TBLR1 monoclonal antibody

Catalog: MB0051

Host: Mouse

Reactivity: Human, Mouse

BackGround:

F-box-like protein involved in the recruitment of the ubiquitin/19S proteasome complex to nuclear receptor-regulated transcription units. Plays an essential role in transcription activation mediated by nuclear receptors. Probably acts as integral component of the N-Cor corepressor complex that mediates the recruitment of the 19S proteasome complex, leading to the subsequent proteasomal degradation of N-Cor complex, thereby allowing cofactor exchange, and transcription activation.

Product:

Purified mouse monoclonal in buffer containing 0.1M Tris-Glycine (pH 7.4, 150 mM NaCl) with 0.2% sodium azide, 50% glycerol

Molecular Weight:

Predicted band size: 60KDa

Observed band size: 60KDa

Swiss-Prot:

Q9BZK7

Purification&Purity:

The antibody was affinity-purified from mouse ascites by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:1000

ICC: 1:50~200

IHC: 1:50~500


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.


Specificity:

This antibody detects endogenous levels of TBLR1 and does not cross-react with related proteins


DATA:



Western blot detection of TBLR1 antibody in Mouse brain, Mouse heart&K562 cell lysates using TBLR1 antibody (1:1000 diluted).



IHC of paraffin-embedded human breast cancer using anti-TBLR1 diluted 1/500-1/1000.



Immunocytochemistry of HeLa cells using anti-TBLR1 antibody diluted 1:150.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151