

**eIF5A monoclonal antibody**

Catalog: MB10470

Host: Rabbit

Reactivity: Human, Mouse, Rat

**BackGround:**

mRNA-binding protein involved in translation elongation. Has an important function at the level of mRNA turnover, probably acting downstream of decapping. Involved in actin dynamics and cell cycle progression, mRNA decay and probably in a pathway involved in stress response and maintenance of cell wall integrity. With syntenin SDCBP, functions as a regulator of p53/TP53 and p53/TP53-dependent apoptosis. Regulates also TNF-alpha-mediated apoptosis.

**Product:**

50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA

**Molecular Weight:**

Calculated MW: 17 kDa; Observed MW: 17 kDa

**Swiss-Prot:**

P63241

**Purification&Purity:**

Affinity Purified

**Applications:**


WB: 1/500-1/1000 IHC: 1/50-1/100 IP: 1/20

**Storage&Stability:**


Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

**Isotype:**

IgG

**DATA:**

Western blot analysis of eIF5A in Jurkat lysates using eIF5A antibody.



Western blot analysis of eIF5A in Jurkat, THP-1, C6, 3T3, Hela lysates using eIF5A antibody.

Immunohistochemistry analysis of paraffin-embedded Human breast cancer using eIF5A antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

**Note:**

For research use only, not for use in diagnostic procedure.

**Bioworld Technology, Inc.**

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: [info@bioworld.com](mailto:info@bioworld.com)

Tel: 6123263284

Fax: 6122933841

**Bioworld technology, co. Ltd.**

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: [info@biogot.com](mailto:info@biogot.com)

Tel: 0086-025-68037686

Fax: 0086-025-68035151